



2157 Royal Windsor Dr.
Mississauga ON

Compounding Specialty
Clarkson Pharmacy

Royal Windsor Drive



Rx@inflex.com

Fax: 1-833-777-0417

Ph: 1-800-701-5262

**FREE SHIPPING
NATIONWIDE**

Patient Name: _____ Phone #: _____

ALOPECIA

- ☐ **HairGain™** Serum X 100mL + 5R
Bimatoprost 0.1% Cetirizine 1% Tretinoin 0.01% Caffeine 0.2% Finasteride 0.25% Minoxidil 7%
Add 6 actuations of HairGain™ serum to provided Applicator Comb. Massage into the scalp for 30 seconds once daily. Do not wash hair for a minimum of 6 hours.
- ☐ **HairGain™** Serum X 100mL + 5R
(Finasteride Free) Bimatoprost 0.1% Cetirizine 1% Tretinoin 0.01% Caffeine 0.2% Minoxidil 7%
Add 6 actuations of HairGain™ serum to provided Applicator Comb. Massage into the scalp for 30 seconds once daily. Do not wash hair for a minimum of 6 hours.

MOLLUSCUM CONTAGIOSUM

- ☐ **MOLLUSCUM CONTAGIOSUM™** X 30mL
FORMULA

Potassium hydroxide 5% Lidocaine 2% Gel Apply BID

Stop applying when pimple appears bright red, scabbed or broken for more than one day. Natural healing of lesions will take place thereafter. Do not use on younger than 2 year old children.

ERECTILE DYSFUNCTION

- ☐ **MAXYM™ S** Sildenafil 100mg X 72 Melt Troches (M:6 Q 14 days). † 15-30 min before coitus. Max 1 Q 24 hrs.
- ☐ **MAXYM™ T** Tadalafil 20mg X 72 Melt Troches (M:6 Q 14 days). † 15-30 min before coitus. Max 1 Q 36 hrs.
- ☐ **MAXYM™ Support** Sildenafil 60mg, Tadalafil 15mg X 72 Melt Troches (M:6 Q 14 days). † 15-30 min before coitus. Max 1 Q36 hrs.
- ☐ **MAXYM™ Peak** Sildenafil 60mg, Tadalafil 18mg, Apomorphine 2mg X 72 Melt Troches (M:6 Q 14 days). † 15-30 min before coitus. Max 1 Q36 hrs
- ☐ **MAXYM™ Ultra** Sildenafil 75mg, Tadalafil 20mg, Apomorphine 2mg X 72 Melt Troches (M:6 Q 14 days). † 15-30 min before coitus. Max 1 Q36 hrs.

Prescriber and patient made an informed decision on the risk and benefit of combining sildenafil and tadalafil in this prescription

PREMATURE EJACULATION

- ☐ **MAXYM™ Duration** Lidocaine 8% Benzocaine 2% spray. 2X10mL. Spray 2-3 sprays on the frenulum of the penis 20 min before coitus.

HYPERPIGMENTATION

- ☐ **DermaBright™** Cream X 30G + 3R. Apply every second day for 2 weeks then once daily thereafter as tolerated.
Hydroquinone 4% Kojic Acid 2% Ascorbic Acid 1% Hydrocortisone 1% Tretinoin 0.025%

PLANTAR WARTS

- ☐ Salicylic Acid 20% Lactic Acid 20% Trichloroacetic Acid 10% Topical Gel X 100 ML + 2R
Apply once daily and occlude with duct tape. Stop after 2-4 weeks of use.

NAIL FUNGUS, Onychomycosis.


- ☐ Ciclopirox 6% Terbinafine 6% Fluconazole 2% Ibuprofen 2% Topical Nail Liquid X 20 ML+ 5R
Apply once daily 6 days a week. Clean with rubbing alcohol once a week. Stop after 4-6 months of use.

MOUTH PAIN

- ☐ Magic Mouth Wash (Benadryl 40mL/Nystatin 40mL/Tetracycline 250mg capX2/Dexamethasone 4 mg/mL 4 mL/Distilled Water 166mL) Swish and expectorate qid.

_____ M.D. CPSO #: _____ Date: _____

Prescriber may cross off or add any ingredient above as she/he see appropriate



Rx@inflex.com
Fax: 1-833-777-0417
Ph: 1-800-701-5262
**FREE SHIPPING
NATIONWIDE**

THE FOLLOWING CREAMS ARE BILLED DIRECTLY TO PATIENT'S INSURANCE BY OUR PHARMACY AND COULD BE READY WITHIN 15-30 MINUTES

Patient Name: _____ Phone #: _____

ARTHRITIS, MILD TO MODERATE MUSCULOSKELETAL CONDITIONS, INFLAMMATORY/INJURY PAIN

- ☐ **InflaFlex** Targeting your source of pain In our proprietary, non-sticky, highly penetrating transdermal delivery emulsion X 200 Gm Pre-packaged Pump.
Ketoprofen 7% Diclofenac 3% Amitriptyline 3% Baclofen 10% Lidocaine 4% Menthol 0.15% Camphor 0.15%

MUSCLE SPASM, SEVERE MUSCULOSKELETAL CONDITIONS

- ☐ **InflaFlex PLUS** Targeting your source of pain In our proprietary, non-sticky, highly penetrating transdermal delivery emulsion X 200 Gm Pre-packaged Pump.
Ketoprofen 7% Baclofen 10% Amitriptyline 3% Cyclobenzaprine 2% Lidocaine 4% Menthol 0.15% Camphor 0.15%

NEUROPATHIC PAIN, CARPAL TUNNEL SYNDROME, TRIGGER FINGER

- ☐ **InflaFlex NEURO** Targeting your source of pain In our proprietary, non-sticky, highly penetrating transdermal delivery emulsion X 200 Gm Pre-packaged Pump.
Ketoprofen 7% Gabapentin 10% Amitriptyline 2% Baclofen 6% Lidocaine 4% Menthol 0.15% Camphor 0.15%

POST-HERPETIC NEURALGIA

- ☐ Amitriptyline 10% Gabapentin 10% Lidocaine 10% Topical Spray
Sig: Spray 1-2 sprays to the affected area 5-10 minutes before applying pain cream Mitte: 100ml
BOTH ☐ **InflaFlex NEURO** Targeting your source of pain In our proprietary, non-sticky, highly penetrating transdermal delivery emulsion X 200 Gm Pre-packaged Pump.
Ketoprofen 7% Gabapentin 10% Amitriptyline 2% Baclofen 6% Lidocaine 4% Menthol 0.15% Camphor 0.15%

FIBROMYALGIA

- ☐ **FibroSyn** Naltrexone 1.5mg cap po od hs to increase by 1.5mg on weekly bases to reach 4.5-6mg /day as tolerated.
Mitte: 945mg dispensed in 1.5,3,4.5 or 6mg caps. Contraindicated for patient currently on opioid(s).

CYSTITIS PAIN

- ☐ Phenazopyridine 200mg Oral Capsules (same active ingredient as in Pyridium®) TID PRN M:21 caps.

DIRECTIONS (all Topical Preparations will have the following directions except if otherwise indicated by prescriber)

Sig: apply 4-6 pumps (2-3 grams) to affected area TID-QID

Mitte: 200 grams (200gm pump = 1 month)

Refills: 5 _____



Ask for the
Original Airless Pump
InflaFlex NEURO
Targeting your source of pain
at no extra charge.



Ask for the
Original Airless Pump
InflaFlex
Targeting your source of pain
at no extra charge.

_____ M.D. CPSO #: _____ Date: _____